

Deeply Virtual Compton Scattering off light nuclei:

a crossroad between hadronic and nuclear Physics toward the 3D nuclear imaging

Sara Fucini

in collaboration with

Sergio Scopetta,

University of Perugia and INFN section of Perugia, Italy,

Michele Viviani

INFN section of Pisa, Italy

Raphaël Dupré,

Université Paris-Saclay, France

February 23, 2021





- **Introduction**
- **(Light) nuclei as a QCD laboratory**
- **Analysis of DVCS off ^4He**
- **Check and test of our models: comparison with JLab data**
- **Future perspectives for the hadronic Physics at the Electron Ion Collider (EIC)**
- **TOPEG: a Monte Carlo event generator for DVCS off light nuclei**

Introduction

Some history

Inclusive DIS process $A(e, e')X$, $Q^2 = -q^2$

$$\frac{d^2\sigma}{d\theta d\nu} \propto F_2^N(x) = \sum e_q^2 x f_q^N(x)$$



- $F_2^N(x)$ is the structure function (observable!)
- $f_q^N(x)$ is the Parton Distribution Function
- $x \equiv x_B = \frac{Q^2}{2P \cdot q} \xrightarrow{\text{LAB frame}} \frac{Q^2}{2M\nu} \xrightarrow{\text{IMF frame}} \text{longitudinal momentum fraction for a quark } q \text{ in a nucleon } N$

In principle $F_2^N = F_2^N(x, Q^2)$: in the Bjorken limit ($\nu, Q^2 \rightarrow \infty$, i.e. x_B fixed), F_2^N scales in x_B

DIS \longrightarrow Incoherent scattering off pointlike partons


The EMC effect

Consider the ratio ($d \approx$ free nucleons)

$$R(x) = \frac{F_2^A(x)}{F_2^d(x)}, \quad x = \frac{Q^2}{2M\nu} \in \left[0; \frac{M_A}{M}\right]$$

The European Muon Collaboration (**EMC**) found $R(x) \neq 1$

The nuclear medium modifies the inner structure of the bound nucleons.




The EMC effect


Consider the ratio ($d \approx$ free nucleons)

$$R(x) = \frac{F_2^A(x)}{F_2^d(x)}, \quad x = \frac{Q^2}{2M\nu} \in \left[0; \frac{M_A}{M}\right]$$

The European Muon Collaboration (**EMC**) found $R(x) \neq 1$



The nuclear medium modifies the inner structure of the bound nucleons.



- $x \leq 0.05$: "Shadowing region"
- $0.3 \leq x \leq 0.85$: "EMC region"
- $0.85 \leq x \leq 1$: "Fermi motion region"

Many models but not yet a complete explanation...

(e.g. see **Cloët et al. JPG (2018)**, for a recent report)

How can we better understand the EMC effect?

- **Elastic scattering** \rightarrow **Form factors** $F(\Delta^2) \rightarrow$ no inner parton structure, only spatial extent




How can we better understand the EMC effect?

- **Elastic scattering** \rightarrow **Form factors** $F(\Delta^2) \rightarrow$ no inner parton structure, only spatial extent
- **Inclusive DIS** \rightarrow **PDFs** $f_q(x, Q^2) \rightarrow$ Longitudinal momentum space, no info on the coordinate plane




How can we better understand the EMC effect?

- **Elastic scattering** \rightarrow **Form factors** $F(\Delta^2) \rightarrow$ no inner parton structure, only spatial extent
- **Inclusive DIS** \rightarrow **PDFs** $f_q(x, Q^2) \rightarrow$ Longitudinal momentum space, no info on the coordinate plane
- **????** $\rightarrow \mathcal{F}_q(x, Q^2, ??..)$ \rightarrow *Transverse coordinate plane and momentum space*



How can we better understand the EMC effect?



- **Elastic scattering** \rightarrow **Form factors** $F(\Delta^2) \rightarrow$ no inner parton structure, only spatial extent
- **Inclusive DIS** \rightarrow **PDFs** $f_q(x, Q^2) \rightarrow$ Longitudinal momentum space, no info on the coordinate plane
- **Exclusive processes** \rightarrow **Generalized parton distributions** $H_q(x, \xi, \Delta^2, Q^2) \rightarrow$ 3-d structure



We can do a *tomography* of hadrons in coordinate space.

Exclusive processes: DVCS off nuclei in handbag approximation

Two different *channels* for DVCS off nuclei: **coherent** and **incoherent**



- Factorization property $\Delta^2 \ll Q^2$ (e.g., see Collins et al., PRD (1997)) :

► **HARD PART** \Rightarrow perturbative QED & QCD

► **SOFT PART** \Rightarrow non-perturbative QCD \rightarrow **GPDs**

- GPDs depend on: $\left(a^\pm = \frac{a_0 \pm a_3}{\sqrt{2}}; \bar{P} = \frac{P+P'}{2} \text{ and } \bar{k} = \frac{k+k'}{2} \right)$

► $\Delta^2 = t = (P' - P)^2 = (q_1 - q_2)^2$

► $\xi = -\frac{\Delta^+}{2\bar{P}^+}$

► $x = \frac{\bar{k}^+}{\bar{P}^+}$

► $Q^2 = -(\kappa - \kappa')^2$

- $x \leq \xi$: GPDs describe **antiquarks**; $-\xi \leq x \leq \xi$: GPDs describe **$q\bar{q}$ pairs**; $x \geq \xi$: GPDs describe **quarks**

GPDs in a nutshell (i)

GPDs are introduced considering the **light-cone correlator**:

$$F_{S,S'}^A = \frac{1}{2} \int \frac{dz^-}{2\pi} e^{ixP^+ z^-} \langle P' S' | \bar{\psi} \left(-\frac{z^-}{2} \right) \gamma^+ \psi \left(\frac{z^-}{2} \right) | P S \rangle$$

$$= \frac{1}{2\bar{P}^+} \left[H_q^A(x, \xi, t) \bar{u}(P', S') \gamma^+ u(P, S) + E_q^A(x, \xi, t) \bar{u}(P', S') \frac{i\sigma^{+\alpha} \Delta_\alpha}{2M} u(P, S) \right]$$

1) Form factor

2) PDFs (when $P = P'$, i.e $t = \xi = 0$)

$$\sum_N \int_{-1}^1 dx \sum_q e_q H_q^A(x, \xi, t) = F_1^A(t)$$


$$H_q^A(x, 0, 0) = q_q^A(x) \quad x > 0$$

$$H_q^A(x, 0, 0) = -\bar{q}_q^A(-x) \quad x < 0$$

3) Probabilistic interpretation in impact parameter space (Burkardt, PRD (2000))

$$\rho^q(x, \vec{b}_\perp) = \int \frac{d^2 \vec{\Delta}_\perp}{(2\pi)^2} e^{-i\vec{b}_\perp \cdot \vec{\Delta}_\perp} H_q^A(x, 0, \Delta_\perp^2)$$


Toward the explanation of the EMC effect



- At JLab kinematics, **Bethe Heitler** process interferes with DVCS enhancing this latter. For this reason, it is convenient to measure **asymmetries**, e.g.

$$A_{LU} = \frac{\sigma^{\lambda=+}_{\sigma^{\lambda=+}} - \sigma^{\lambda=-}_{\sigma^{\lambda=-}}}{\sigma^{\lambda=+}_{\sigma^{\lambda=+}} + \sigma^{\lambda=-}_{\sigma^{\lambda=-}}}$$

$$\sigma^{\lambda} \propto T_{BH}^2 + T_{DVCS}^2 + \mathcal{I}_{BH-DVCS}^{\lambda}$$



that can be expressed in terms of

- Form Factors**

$$T_{BH} \propto FF(\Delta^2)$$

- Compton Form Factors** ($CCFs \propto$ GPDs)

$$T_{DVCS} \propto \mathcal{H}(\xi, \Delta^2) = \int_{-1}^1 dx \frac{H_q^A(x, \xi, \Delta^2)}{x - \xi + i\epsilon} = \Re \mathcal{H}(\xi, \Delta^2) + i \Im \mathcal{H}(\xi, \Delta^2)$$



To de-convolute GPDs we need a wide range in t and ξ otherwise a model input is needed \rightarrow first results for proton's tomography **Dupré et al. PRD (2017) 95, p. 01150**

- DVCS is sensitive only to chiral-even GPDs and is dominated by quark GPDs in the valence region.

Why nuclei?

Coherent DVCS channel

Handbag approximation



Impulse approximation (IA)

Impulse approximation in light nuclei

As a good starting point, let us consider the **IA** (in a second step we can add as many refined ingredients as we want) whose validity has been experimentally confirmed at JLab kinematics in, e.g., **Slifer et al. PRL (2008) 022303**.

Let us start from the light-cone correlator where we insert two complete sets of states

- the active nucleon (kinematically off-shell)
- the remnant $A - 1$ -body system

We get a **convolution formula for the GPD** $H_q^A(x, \xi, \Delta^2)$, $\left(z = \frac{\text{long. } N \text{ mom. fraction}}{\text{long. } q \text{ mom. fraction}}\right)$

$$H_q^A(x, \xi, \Delta^2) \approx \sum_N \int \frac{dz}{z} h_N^A(z, \xi, \Delta^2) H_q^N\left(\frac{x}{z}, \frac{\xi}{z}, \Delta^2\right)$$

where the **off-diagonal light-cone momentum distribution** is

$$h_N^A(z, \xi, \Delta^2) = \int dE d\vec{p} P_N^A(\vec{p}, \vec{p} + \vec{\Delta}, E) \delta\left(z - \frac{\bar{p}^+}{\bar{P}^+}\right)$$

$P_N^A(\vec{p}, \vec{p} + \vec{\Delta}, E)$ is the **one body off-diagonal spectral function** of the nucleon N in the nucleus A

Impulse approximation in light nuclei (ii)


The **forward limit** (i.e. $\xi, \Delta^2 \rightarrow 0$) of the **light-cone momentum distribution**

$$h_N^A(z, 0, 0) = f_N^A(\tilde{z}) = \int dE \int d\vec{p} P_N^A(\vec{p}, E) \delta\left(\tilde{z} - \frac{p^+}{P^+}\right)$$

and the forward limit of the GPD becomes:

$$H_q^A(x, 0, 0) = q_q^A(x) \approx \sum_N \int_1^x \frac{d\tilde{z}}{\tilde{z}} f_N^A(\tilde{z}) q_q^N\left(\frac{x}{\tilde{z}}\right)$$

where $f_N^A(\tilde{z})$ strongly peaked around $\tilde{z} \approx 1/A$



For ${}^4\text{He}$, $f_N^A(z)$ picked at $z \approx 0.25$

How can the nuclear effects be inferred from $f_N^A(z)$?

ξ is the fraction of "+" momentum transfer and cannot exceed the width of $f_N^A(z)$ to have the target intact after the interaction.

If DVCS were observed in a wide range of t (ξ), exotic effects beyond IA, e.g. non-nucleonic d.o.f., would be pointed out (**Berger et al. PRL 87 (2001)**).

Similar effect predicted in DIS at $x_B > 1$, where data are not accurate enough

When dealing with **nuclear targets**, keep in mind that:

- a system of spin S has $(2S + 1)^2$ parton helicity conserving and chiral-even quark GPDs and $(2S + 1)^2$ parton helicity flipping and chiral odd quark GPDs
 $\Rightarrow 2(2S + 1)^2$ GPDs. Considering NLO terms, we have $4(2S + 1)^2$ **GPDs**.
- for **light nuclei**, realistic calculations of the wave functions, exact solutions of the Schrödinger equation with **phenomenological NN potentials** (e.g. Av18) and 3-body forces, are possible

When dealing with **nuclear targets**, keep in mind that:

- a system of spin S has $(2S + 1)^2$ parton helicity conserving and chiral-even quark GPDs and $(2S + 1)^2$ parton helicity flipping and chiral odd quark GPDs
 $\Rightarrow 2(2S + 1)^2$ GPDs. Considering NLO terms, we have $4(2S + 1)^2$ **GPDs**.
- for **light nuclei**, realistic calculations of the wave functions, exact solutions of the Schrödinger equation with **phenomenological NN potentials** (e.g. Av18) and 3-body forces, are possible

Theoretical challenge: the bigger is A , the harder are such calculations

- **Deuteron** ($S = 1$): many GPDs yet at leading twist (**Berger et al. PRL (2001)**) evaluated in a light front framework **Cano et al. EPJA (2003)**) also in the transversity sector (**Cosyn et al. PRD (2018)**), access to the neutron in the unpolarized setup of the incoherent channel.
- **^3He** ($S = 1/2$): study of the isospin-flavor dependence of nuclear effects (**Scopetta PRC (2004)**, **Scopetta PRC (2009)**), evaluation of its conventional nuclear structure (e.g. **Rinaldi et al. PRC (2012)**, **Rinaldi et al. FBS (2014)**); not yet DVCS data for a ^3He target; preliminary results for the observables in **Fucini et al. FBS (2021)**

Why is ${}^4\text{He}$ a golden nucleus?

- ${}^4\text{He}$ is a typical few body system and it is theoretically well known
- exact and realistic calculations are difficult BUT possible
- $J_{4He}^{\pi} = 0^{+}$ and $I_{4He} = 0 \implies$ only one chiral-even GPD at LO
- CLAS and ALERT collaboration are carrying on an experimental program at JLab using ${}^4\text{He}$ target

Coherent (**PRL 119, 202004 (2017)**) and incoherent (**PRL 123, 032502 (2019)**) DVCS off ${}^4\text{He}$ has been measured at the Jefferson Laboratory!

- good perspectives at **JLab with a 12 GeV** electron beam and the forthcoming **EIC**

Our point is to obtain models able to distinguish “conventional” and “exotic” nuclear structure effects

Coherent DVCS off ^4He

A convolution formula for the chiral even GPD H_q can be obtained in terms of:

- GPDs of the inner nucleons**

$$H_q^{4He}(x, \xi, \Delta^2) = \sum_N \int_{|x|}^1 \frac{dz}{z} h_N^{4He}(z, \xi, \Delta^2) \mathbf{H}_q^N\left(\frac{x}{\xi}, \frac{\xi}{\xi}, \Delta^2\right)$$

- light-cone momentum distribution**

$$h_N^{4He}(z, \Delta^2, \xi) = \frac{M_A}{M} \int dE \int_{p_{min}}^{\infty} dp \int_0^{2\pi} d\phi p \tilde{M} P_N^{4He}(\vec{p}, \vec{p} + \vec{\Delta}, E)$$

$$\xi_A = \frac{M_A}{M} \xi, \tilde{z} = z + \xi_A,$$

$$\tilde{M} = \frac{M}{M_A} \left(M_A + \frac{\Delta^+}{\sqrt{2}} \right), p_{min} = f(z, \xi_A, E), \mathbf{H}_q^N = \sqrt{1 - \xi^2} [H_q^N - \frac{\xi^2}{1 - \xi^2} E_q^N]$$

One needs the **non-diagonal spectral function** and the **nucleonic GPDs** (we used the Goloskokov-Kroll models (EPJ C (2008)- EPJ C (2009))

The ^4He spectral function: off diagonal case


$$P_N^{4He}(\vec{p}, \vec{p} + \vec{\Delta}, E) = \rho(E) \sum_{\alpha \sigma} \langle P + \Delta | -p \alpha, p + \Delta \sigma \rangle \langle p \sigma_N, -p \alpha | P \rangle$$

with **removal energy** $E = |E_A| - |E_{A-1}| - E^*$



Ground-state contributions

2-body channels

- $\langle ^4\text{He} | p, ^3\text{H} \rangle$;
- $\langle ^4\text{He} | n, ^3\text{He} \rangle$;



Excited-state contributions



3-body channels

- $\langle ^4\text{He} | p, d n \rangle$;
- $\langle ^4\text{He} | n, d p \rangle$;

4-body channels

- $\langle ^4\text{He} | n, p n p \rangle$;
- $\langle ^4\text{He} | p, n p n \rangle$.

$$\begin{aligned}
 P_N^{4He}(\vec{p}, \vec{p} + \vec{\Delta}, E) &= n_0(\vec{p}, \vec{p} + \vec{\Delta})\delta(E) + P_1(\vec{p}, \vec{p} + \vec{\Delta}, E) \\
 &= n_0(|\vec{p}|, |\vec{p} + \vec{\Delta}|, \cos\theta_{\vec{p}, \vec{p} + \vec{\Delta}})\delta(E) + P_1(|\vec{p}|, |\vec{p} + \vec{\Delta}|, \cos\theta_{\vec{p}, \vec{p} + \vec{\Delta}}, E) \\
 &\simeq a_0(|\vec{p}|)a_0(|\vec{p} + \vec{\Delta}|)\delta(E) + \sqrt{n_1(|\vec{p}|)n_1(|\vec{p} + \vec{\Delta}|)}\delta(E - \bar{E})
 \end{aligned}$$

- the **total momentum distribution** is $n(p)$

$$n_1(|\vec{p}|) = n(|\vec{p}|) - n_0(|\vec{p}|)$$

- $n_0(k)$ is the momentum distribution when the recoiling system in the **ground-state**

$$n_0(|\vec{p}|) = |a_0(|\vec{p}|)|^2$$

with


$$a_0(|\vec{p}|) = \langle \Phi_3(1, 2, 3) \chi_4 \eta_4 | j_0(|\vec{p}| R_{123,4}) \Phi_4(1, 2, 3, 4) \rangle .$$

- $n(p)$ has been evaluated for the 4-body and 3-body systems within the **Av18 NN interaction (Wiringa et al., PRC (1995)) + UIX 3-body forces (Pudliner et al., PRL (1995))**
- \bar{E} is the **average excitation energy** of the recoiling system (the model for the excited part of the diagonal s.f. **M. Viviani et al., PRC (2003)** is a realistic update of the model of **Ciofi et al., PRC (1996)**, i.e. $P_N^{1 \text{ our model}} = N(p)P_N^{1 \text{ Ciofi's model}}$)

$$\Im \mathcal{H}_A(\xi, t) = \sum_{q=u,d,s} e_q^2 (H_q^A(\xi, \xi, \Delta^2) - H_q^A(-\xi, \xi, \Delta^2))$$

$$\Re \mathcal{H}_A(\xi, t) = \text{Pr} \sum_{q=u,d,s} e_q^2 \int_0^1 \left(\frac{1}{\xi+x} + \frac{1}{x-\xi} \right) (H_q^A(x, \xi, t) - H_q^A(-x, \xi, t))$$

Black squares \rightarrow JLab data from the CLAS coll. (**Hattawy et al., PRL (2018)**)




Beam spin asymmetry as a function of azimuthal angle $\phi = 90^\circ$

$$A_{LU}(\phi) = \frac{\alpha_0(\phi) \Im m(\mathcal{H}_A)}{\alpha_1(\phi) + \alpha_2(\phi) \Re e(\mathcal{H}_A) + \alpha_3(\phi) \left(\Re e(\mathcal{H}_A)^2 + \Im m(\mathcal{H}_A)^2 \right)}.$$


where $\alpha_i(\phi)$ are kinematical coefficients from **A. V. Belitsky et al., Phys. Rev. D 79, 014017 (2009).**

From left to right, the quantity is shown in the experimental Q^2 , x_B and $-t$ bins

Results of our approach VS **EG6 data**



Incoherent DVCS off ^4He



The **beam spin asymmetry** (BSA) measured is:

$$A_{LU} = \frac{d\sigma^+ - d\sigma^-}{d\sigma^+ + d\sigma^-}$$

where

\pm refers to positive(negative) beam polarizations.

Fundamental starting points for our **Impulse Approximation** approach are:

- kinematical **off shellness**:

$$p_0 = M_A - \sqrt{M_{A-1}^{*2} + \vec{p}^2} \simeq M_N - E - T_{rec} \implies p^2 \neq m^2$$


- general expression for **cross section**

$$(d\sigma^\pm)_{INC} = (2\pi)^4 \frac{1}{2P_A \cdot k} \sum_N \sum_X |\mathcal{A}^\pm|^2 \delta^4(P_A + k - k' - p_X - p_N - q_2) LIPS$$

where $LIPS = d\tilde{p}_X d\tilde{k}' d\tilde{q}_2 d\tilde{p}_N$

Our formalism (i)

In a frame where the target nucleus is at rest, the cross section and its azimuthal dependence are expressed in terms of a **convolution formula** between:



- the **diagonal** spectral function

$$d\sigma_{Incoh}^{\pm} = \int_{exp} dE d\vec{p} \frac{p \cdot k}{p_0 |\vec{k}|} P^{4He}(\vec{p}, E) d\sigma_b^{\pm}(\vec{p}, E, K)$$

- the DVCS cross section off a bound proton

The differential cross section appearing in A_{LU} is


$$\frac{d\sigma_{Incoh}^{\pm}}{dx_B dQ^2 d\Delta^2 d\phi} = \int_{exp} dE d\vec{p} P^{4He}(\vec{p}, E) |\mathcal{A}^{\pm}(\vec{p}, E, K)|^2 g(\vec{p}, E, K)$$

where

- $K = \{x_B = \frac{Q^2}{2M\nu}, Q^2, \phi, \Delta^2\}$ fixes the proper range of integration
- $g(\vec{p}, E, K)$ arises from the integration of LIPS and includes also the flux factor

Our formalism (ii)

Schematically $d\sigma^\pm \approx \int d\vec{p} dE P^{4He}(\vec{p}, E) |A^\pm(\vec{p}, E, K)|^2$ with
 $|A^\pm|^2 = \mathcal{T}_{BH}^2 + \mathcal{T}_{DVCS}^2 + \mathcal{I}_{DVCS-BH}^\pm$.



The BSA for the incoherent DVCS reads:

$$A_{LU}^{Incoh}(K) = \frac{\mathcal{I}^{4He}(K)}{\mathcal{T}_{BH}^{24He}(K)}$$

$$\mathcal{I}^{4He}(K) = \int_{exp} dE d\vec{p} P^{4He}(\vec{p}, E) g(\vec{p}, E, K) \mathcal{I}(\vec{p}, E, K)$$

$$\mathcal{T}_{BH}^{24He}(K) = \int_{exp} dE d\vec{p} P^{4He}(\vec{p}, E) g(\vec{p}, E, K) \mathcal{T}_{BH}^2(\vec{p}, E, K)$$


$$A_{LU}^{Incoh}(K) = \frac{\mathcal{I}^{4He}(K)}{T_{BH}^{24He}(K)}$$

- Our expression for $|\mathcal{T}_{BH}(\vec{p}, E, K)|^2 = c_0^{BH} + c_1^{BH} \cos(\phi) + c_2^{BH} \cos(2\phi)$ is a generalization for a moving bound nucleon of results by **Muller et al., NLB (2002)**
- the interference BH-DVCS $\mathcal{I}(\vec{p}, E, K) \approx s_1^{\mathcal{I}}(\vec{p}, E, K) \Im m \mathcal{H}(\xi', \Delta^2, Q^2)$.
- For the proton GPD H_q^N , again, we used **GK model** evaluated for $\xi' = \frac{Q^2}{(p+p_N)(q_1+q_2)} \neq \frac{x_B}{2-x_B} = \xi_{rest}$
- No nuclear modifications occur for the **form factors** of the bound proton
- For the diagonal spectral function $P^{4He}(\vec{p}, E)$ we use an Av18-based model (**M. Viviani et al., PRC 67, 034003 (2003)**)
 - the **ground-state** of the recoiling system is described in terms of exact wave functions for the 4-body and 3-body systems
 - the **excited state** of the recoiling system is an update of the 2-nucleon correlation model by **Ciofi et al., PRC 53 1689 (1996)**.

Incoherent DVCS: results

- Our results are compared with the experimental data from EG6 collaboration at JLab (**M. Hattawy et al., PRL 123, 032502 (2019)**).

From left to right, the quantity is shown in the experimental Q^2 , x_B and $-t$ bins




✓ Good agreement in the region of high Q^2

An analysis of the interplay between the t and Q^2 dependence could reveal if FSI effects could be responsible of the disagreement in low Q^2 region

Testing the IA

Let us consider the **MMS13 model** (Mezrag et al., PRD (2013)) for the proton GPD and the **closure approximation** (fixed removal energy \Rightarrow momentum distribution)




Our BSA turns out to be :

- sensitive to the nucleonic model used, in particular at low values of Q^2
- mildly sensitive to the details of the nuclear model used in the calculation


Nuclear dynamics in $|\text{BH}|^2$ and in the BH-DVCS interference

Are the nuclear effects measured depending on the modification of the bound proton partonic structure? Let us consider the ratio

$$A_{LU}^{Incoh}/A_{LU}^p = \frac{\mathcal{I}^{4He}}{\mathcal{I}^p} \frac{T_{BH}^{2p}}{T_{BH}^{2^{4He}}} = \frac{R_I}{R_{BH}} \propto \frac{(nucl.eff.)_{\mathcal{I}}}{(nucl.eff.)_{BH}},$$



Using the GK models and the spectral function




Nuclear dynamics modify both $|T_{BH}|^2$ and $\mathcal{I}_{BH-DVCS}$ but in the ratio these effects **compensate** each other

This fact hasn't to do with a modification of the parton structure

as confirmed by:

- the ratio A_{LU}^{Incoh}/A_{LU}^p for "pointlike" protons
- the "EMC-like" trend

$$R_{EMC-like} = \frac{1}{\mathcal{N}} \frac{\int_{exp} dE d\vec{p} P^4 He(\vec{p}, E) \Im m \mathcal{H}(\xi', \Delta^2)}{\Im m \mathcal{H}(\xi, \Delta^2)}$$



DVCS off light nuclei at the EIC


- ▶ **JLab** (fixed target experiments) \rightarrow mostly x_B valence region
- ▶ A worldwide community (+ 1000 users) and a strong activity for a global project: the **Electron Ion Collider** (EIC)

A groundbreaking boost in our knowledge at the EIC in the next decades where **DVCS is a key process** (exclusive reactions WG)

- Large range of center-of-mass energies and very high **luminosity** \rightarrow high precision measurements
- **Polarization of nuclear beams** \rightarrow spin asymmetries (e.g ^3He can be actually used as a neutron target)
- ... not only nuclear tomography
 - **Role of gluon GPDs and shadowing effects**: possible gluon d.o.f. in nuclei will be accessible at very small values of x_B (**Goeke et al. PRC (2009)**)
 - Study of the **nuclear energy momentum tensor** and the **distribution of pressure and shear forces** inside the nucleus (**M.V. Polyakov, PLB (2003)**): from the energy-momentum tensor, the total angular momentum of the target can be accessed.

TOPEG: a Monte Carlo event generator for DVCS off light nuclei

TOPEG (The Orsay-Perugia Event Generator) is a `Root` based generator (S. Jadach (2005)) + **our model** for the coherent DVCS



+

Our model for
the coherent
DVCS
*Fucini et al., PRC
98 (2018)*



TOPEG
(The Perugia-
Orsay Event
Generator)

- ▶ Check for JLab 6 GeV
- ▶ We generated events for the three energy configurations for the DVCS off ^4He at the EIC
 - 5x41 GeV
 - 10x110 GeV
 - 18x110 GeV
- ▶ These results will be included in the **Yellow Report of the EIC user group** (to be released by the end of this month)




18 x 110 GeV: kinematical distributions

We generated events weighted by the cross section $\frac{d^4\sigma}{dQ^2 dt d\phi dx_B}$

- 1 million events
- in the x-section, we set $\Re e(\text{CFF})=0$ (limitation in the computation time)
- Luminosity: 250 nb^{-1} (**NOT ENOUGH!!**)
- $Q^2 > 2 \text{ GeV}^2$, $y < 0.8$, $t_{\min} < |t| < t_{\min} + 0.5 \text{ GeV}^2$

For small $|t|$, we expect an enhancement of the cross section for the dominance of the BH process ($\simeq \text{FF}^2$).

$$|t_{\min}| = \frac{4M_{4H_e}^2 \xi^2}{1 - \xi^2} \quad \text{with} \quad \xi = \frac{x_B}{2 - x_B} \quad \text{and} \quad x_B = \frac{Q^2}{y(s - M_{4H_e}^2)}$$






Is there plenty of room to study the region around the first diffraction minimum in the ^4He FF ($t_{\text{dif. min}} = -0.48 \text{ GeV}^2$)?

- 99%+ electrons and photons are in the acceptance of the detector matrix
- This is true for all energy configurations

Electrons and photons appear in easily accessible kinematics according to the detector matrix requirements (exceptions for small angles photons)

- Acceptance at low $-t$ will be cut passing through the detectors
 - ▶ t_{min} is set by the detector features (i.e. the Roman pots capabilities)
 - ▶ t_{max} is fixed by the luminosity (billion of events to generate)



Conclusions and outlooks

Our workable approaches to DVCS off ^4He allow to constrain conventional nuclear effects.

● Formal development of a theoretical formula for the chiral even **GPD** of the ^4He with an overall good agreement with JLab data

● Calculation of the **beam spin asymmetry of a bound proton and study of the nuclear effects**

● Concerning the nuclear ingredient, to date we have a s.f.:

- Realistic AV18 + UIX momentum dependence
- Dependence on E , angles and Δ in the s.f. is modeled and not yet realistic

● **Version 1.0 of TOPEG** for a key process at the EIC to make predictions about the cross sections, contributing to the **physical program** and the **design** of the EIC

Our approach is helpful for planning new measurements, not only for interpreting the present data.


- Evaluation of the incoherent channel considering **Final State Interaction** effects (tagged experiments by **ALERT collaboration**)
- **Need for covariance**: relativistic description for both DVCS channels in a **light-front scenario** to achieve polynomiality for GPDs and sum rules in DIS (in our approach, number of particles and momentum sum rule not fulfilled at the same time)
- A full realistic evaluation of the **(off)-diagonal spectral function**
- Concerning **TOPEG**:
 - Preliminary results for the **projections for the ^4He profiles**
 - Study the impact of **non nucleonic d.o.f.**: is this Physics accessible at the EIC ?
 - Re-do the simulations for $\Re e(\text{CFF}) \neq 0$ (tech improvements and possible parallelization to shorten the calculation time and get higher luminosity)
 - Include **shadowing effects** at low x_B
 - Plug-in **other light nuclei targets** and the proton (**free** and **bound**) target
 - Add other hard processes?

Thank you ...
Questions?

Backup slides

Form factor of the ${}^4\text{He}$ at high Q^2

Red dashed line: One body part of the form factor from a direct integration of the diagonal momentum distribution of the ${}^4\text{He}$ within Av18+UIX calculation (figure from **Phys. Rev. Lett. 112, 132503**)



EMC effect with our model for the off diagonal spectral function


$$R(x) = \frac{F_2^{4He}(x)}{F_2^d(x)} \quad x \in [0 : M_A/M]$$

where the **function structures** F_2 for $A = {}^4\text{He}, d$ are defined as

$$F_2^A(x) = \sum_N \int_x^{M_A/M} dz f_N^A(z) F_2^N\left(\frac{x}{z}, Q^2\right)$$



in terms of the *light-cone momentum distribution*

$$f_N^A(z) = \int d\vec{p} \int dE P_N^A(\vec{p}, E) \frac{p^+}{p_0} \delta\left(z - \sqrt{2} \frac{p^+}{M_A}\right)$$



- Our model isn't predictive at **small x**
- Good agreement in the **valence region**
- Strong dependence on the model for F_2^N at **large x**
- Need to better unravel the Q^2 dependence of $R(x)$
Data from **Seely et al., PRL (2009)**

Some checks for our model for the coherent DVCS off ^4He



- **EMC-like effect**

$$R_q(x, 0, 0) = \frac{H_q^A(x_A, 0, 0)}{2(H_q^p(x_A, 0, 0) + H_q^n(x_A, 0, 0))}$$

✓ Good EMC-like behavior;

- **Charge form factor**



$$F_C^{4He}(\Delta^2) = \frac{1}{2} \sum_q e_q \int_0^1 dx H_q^{4He}(x, \xi, \Delta^2)$$

Data (●) from **PRC 160, 4 (1987)**,
theoretical one-body calculation (▲) by
Marcucci et al., PRC 58, 3069 (1998).

✓ Good agreement with the experimental data.

Why choose the treatment with the spectral function

The effects in the numerator and in the denominator of A_{LU}^{Incoh} compensate each other in the ratio. In the **closure approximation**,



- the **removal energy** is fixed to an average value
- the **change of the off-shellness** of the proton produce a big effect in each amplitude

If the nuclear dynamics modifies the amplitudes, the effect can be big even if the parton structure of the bound proton doesn't change